Mon. Not. R. Astron. Soc. **000**, 1–8 (1999)   Printed 5 February 2008   (MN LaTeX style file v1.4)

arXiv:astro-ph/9806378v2 9 Apr 1999

Secular Instability of g-Modes in Rotating Neutron Stars

Dong Lai
Centre for Radiophysics and Space Research, Department of Astronomy, Cornell University, Ithaca, NY 14853, USA
E-mail: dong@spacenet.tn.cornell.edu

Accepted 1999,4; Received 1998 xxx; in original form 1998 xxx

ABSTRACT

Gravitational radiation tends to drive gravity modes in rotating neutron stars unstable. For an inviscid star, the instability sets in when the rotation frequency is about 0.7 times the corresponding mode frequency of the nonrotating star. Neutron stars with spin frequencies $\gtrsim 100$ Hz are susceptible to this instability, with growth time of order years. However, it is likely that viscous dissipation suppresses the instability except for a narrow range of temperatures around 10^9 K. We also show that the viscosity driven instability of g-modes is absent.

Key words: stars: neutron – stars: oscillation – stars: rotation – hydrodynamics – gravitation

1 INTRODUCTION

It is well known that globally nonaxisymmetric instabilities can develop in rapidly rotating neutron stars. Of particular interest is the secular instability driven by gravitational radiation, as first discovered by Chandrasekhar (1970) for the bar-mode ($m = 2$) of an incompressible Maclaurin spheroid, and later shown by Friedman & Schutz (1978a,b) to be a generic feature of rotating stars. The Chandrasekhar-Friedman-Schutz instability (hereafter CFS instability) occurs whenever a backward-going mode (with respect to the rotation) in the corotating frame of the star is "dragged" by rotation to become forward-going in the inertial frame. A CFS unstable mode has negative energy (in the inertial frame), and gravitational radiation makes the energy even more negative.

The CFS instability has been extensively studied for f-modes since it is widely believed that the maximum angular velocities of neutron stars are determined by instabilities of these modes (e.g., Ipser & Lindblom 1990, 1991; Cutler & Lindblom 1992; Lindblom 1995; Yoshida & Eriguchi 1995; Stergioulas & Friedman 1998). The f-mode instabilities occur at rotation frequencies comparable to the maximum "break-up" frequency (~ 1000 Hertz). It was found that viscous dissipation in the neutron star stablizes almost all f-modes except in certain limited temperature regimes. Recently Andersson (1998) pointed out that r-modes (previously studied by Papaloizou & Pringle 1978, Provost et al. 1981, Saio 1982, etc.) are also destablized by gravitational radiation (see also Friedman & Morsink 1998). Moreover, the couplings between gravitational radiation and r-modes (mainly through current multipole moments) are so strong that viscous forces present in hot young neutron stars are not sufficient to suppress the instability (Lindblom et al. 1998; Andersson et al. 1998a). It is therefore likely that gravitational radiation can carry away most of the angular momentum of a nascent neutron star through the unstable r-modes, and the gravitational waves generated in this process could potentially be detectable by LIGO (Owen et al. 1998). The implications of the r-mode instability for accreting neutron stars have also been discussed (Andersson et al. 1998b; Bildsten 1998; Levin 1998).

A neutron star also possesses p-modes and g-modes, whose stability properties have not been studied. The p-modes are similar to the f-modes, except that, as compared to the f-modes, they have higher frequencies, weaker couplings to the gravitational radiation and stronger viscous dampings. Thus the p-mode instabilities, if exist, are certainly less important than the f-mode instabilities. The situation with g-modes is not clear, however. Gravity modes in neutron stars arise from composition (proton to neutron ratio) gradient in the stellar core (Reisenegger & Goldreich 1992; Lai 1994), density discontinuities in the crust (Finn 1987; Strohmayer 1993) as well as thermal buoyancy associated with finite temperatures, either due to internal heat (McDermott, Van Horn & Hansen 1988) or due to accretion (McDermott & Taam 1987; Bildsten & Cutler 1995; Strohmayer & Lee 1996; Bildsten & Cumming 1998). For a nonrotating neutron star, the typical frequencies of low-order g-modes are around 100 Hz and are smaller for higher-order modes. Such low frequencies imply that, contrary to f-modes, g-modes may be destablized by gravitational wave when the star rotates at a rate much smaller than the break-up spin frequency. Moreover, since g-modes possess small, but nonzero mass quadrupole moments, it is not clear *a priori* whether the driving due to gravitational radiation is stronger or weaker for g-modes than that for r-modes (which have zero mass quadrupole moment). The purpose of this

© 1999 RAS

paper is to address these issues related to the secular instability of g-modes.

Our paper is organized as follows. We start in Section 2 by summarizing the essential properties of g-modes in nonrotating neutron stars and defining notations. In Section 3 we calculate the effect of rotation on the mode frequencies and determine the onset of CFS instabilities of g-modes for inviscid stars. In Section 4 we evaluate the driving/damping rates of the g-modes due to gravitational radiation and viscosities. Section 5 considers the possibility of viscosity-driven g-mode instability. In Section 6 we compare the instabilities of g-modes and other modes (f and r-modes) and discuss some possible astrophysical applications of our results.

2 PREPARATIONS: CORE G-MODES OF NEUTRON STARS

In this paper, we shall focus on the core g-modes, with buoyancy provided by the gradient of proton to neutron ratio in the neutron star interior (Reisenegger & Goldreich 1992). These modes exist even in zero-temperature stars. The other types of g-modes (due to crustal density discontinuities or thermal entropy gradient) either have weaker restoring forces or are confined to the outer region of the star and therefore are less susceptible to the CFS instabilities as compared to the core g-modes (see below).

We construct equilibrium neutron star models by solving Newtonian hydrostatic equations. Centrifugal distortion to the stellar structure is neglected as we shall only consider stars with rotation rates far below the break-up value. The equations of state (EOS) used for our neutron star models and g-mode calculations have been described in Lai (1994). The nuclear symmetry energy plays an important role in determining the proton fraction $x = n_p/n$ (where n is the baryon number density and n_p in the proton number density) and hence the frequencies of g-modes. Stable stratification of the liquid core (thus the existence of core g-modes) arises from the x-gradient, and from the fact the β-equilibration time is much longer than the dynamical times of the modes. The Brunt-Väisälä (angular) frequency N is given by

$$N^2 = -\frac{g}{\rho}\left(\frac{\partial\rho}{\partial x}\right)_P \left(\frac{dx}{dr}\right), \tag{1}$$

where $g > 0$ is the local gravitational accelaration. As an estimate of the mode frequency, we neglect the nucleon interaction, and find $x \simeq 5.6\times10^{-3}(\rho/\rho_{\rm nuc})$, where $\rho_{\rm nuc} = 2.8\times10^{14}$ g cm^{-3} is the nuclear density. The Brunt-Väisälä frequency is then

$$N = g\left(\frac{3\rho}{10P}x\right)^{1/2} \sim (G\rho x)^{1/2} \sim 10^3\ {\rm s}^{-1}. \tag{2}$$

Thus the core g-modes typically have frequencies less than $100 - 200$ Hz.

In the following sections, we shall consider two representative models based on a parametrized EOS derived from Wiringa, Fiks & Fabrocini (1988), and labeled as UU and AU in Lai (1994). Figure 1 depicts the profiles of the Brunt-Väisälä frequency, $\nu_N = N/(2\pi)$, as a function of radius inside the star for the two models. We see that $\nu_N = 0$ at the center (where $g = 0$) and increases outward because of decreasing sound speed c_s and increasing g (despite that x decreases outward). We have set $\nu_N = 0$ outside the core ($\rho \lesssim 10^{14}$ g cm^{-3})[★], and thus eliminated the crustal modes (see below). For completeness, Figure 1 also shows the Lamb frequency (or the acoustic cut-off frequency), $\nu_L = \sqrt{l(l+1)}\,c_s/(2\pi r)$, with $l = 2$. Gravity wave (with frequency ν) can propagate only in the region satisfying $\nu < \nu_L$ and $\nu < \nu_N$. For a typical $\nu \sim 100$ Hz, the wave propagation region lies in the range $0.5 \lesssim r/R \lesssim 0.9$. For convenience, the basic properties of the modes for nonrotating stars are summarized in Table 1.



Figure 1. Profiles of the Brunt-Väisälä frequency ν_N (heavy curves) and the Lamb frequency ν_L (light curves) for two neutron star models: UU (solid curves) and AU (dashed curves). We have set $\nu_N = 0$ outside the core ($\rho \lesssim 10^{14}$ g cm^{-3}). The dips in the ν_L curves at $r/R \simeq 0.94$ correspond to neutron drip at $\rho \simeq 4\times10^{11}$ g cm^{-3}.

The crustal discontinuity modes have angular frequencies approximately given by $\omega \sim (g/R)^{1/2}(\Delta\rho/\rho)^{1/2}$. With typical density discontinuity $\Delta\rho/\rho \sim 1\%$, this gives a frequency of order 200 Hz (Finn 1987). Although these modes can have frequencies greater than the core g-modes, the crustal modes are confined in the outer region of the star and therefore have very small mass quadrupole moments[†]. The thermally driven g-modes (McDermott et al. 1988) have frequencies less than 50 Hz for $T \lesssim 10^{10}$ K, indicating smaller restoring forces as compared to the core g-modes. We will not consider the crustal modes and thermal modes in the rest of this paper.

[★] The transition from the core (where only free nucleons and electrons exist) to the inner crust (where heavy nuclei co-exist with free neutrons) is somewhat ambiguous. The inner crust is described by the equation of state of Baym, Bethe & Pethick (1971) while the core is described by that of Wiringa et al. (1988). We have found the g-mode properties to be insensitive to the actual transition radius within the allowed range.

[†] This is indicated by the fact that the mode damping time due to gravitational radiation is rather long, exceeding $10^4 - 10^5$ years.

Table 1. Core g-modes of nonrotating neutron stars

Model	M $(M_\odot)$	R (km)	ρ_c (g cm^{-3})	mode [a]	ν (Hz)	$\lvert\delta D_{22}\rvert$ [b]
UU	1.4	13.48	6.95×10^{14}	g_1	148	6.2×10^{-4}
				g_2	97	1.8×10^{-5}
AU	1.4	12.37	8.18×10^{14}	g_1	72	1.1×10^{-4}
				g_2	52	9.9×10^{-6}

[a] The subscript n in g_n specifies the radial order of the mode. The angular order is $l = 2$.

[b] Note that the values of $|\delta D_{22}|$ quoted here differ from those in Lai (1994) by up to 80%. This is because we adopt Cowling approximation in this paper while in Lai (1994) the full equations are solved; see discussion at the end of §3.

3 ONSET OF INSTABILITY: INVISCID STARS

For neutron stars consisting of ideal fluid (neglecting viscosity), it is possible to determine the onset of CFS instability of g-modes without calculating the growth/damping rates due to dissipative processes. Consider a mode with azimuthal angular dependence $e^{im\phi}$ and time dependence $e^{i\omega_r t}$, where ω_r (> 0) is the mode angular frequency in the rotating frame. The rotation angular frequency is Ω_s (> 0). A mode with $m > 0$ propagates in the direction opposite to the rotation, with pattern speed $(-\omega_r/m)$ (in the rotating frame). In the inertial frame, the mode frequency is

$$\omega_i = \omega_r - m\Omega_s. \tag{3}$$

When $\omega_i < 0$, the mode attains a prograde pattern speed, $-(\omega_i/m)$, in the inertial frame, and the CFS instability sets in (Friedman & Schutz 1978a,b).

Since we shall consider Ω_s comparable to ω_r, a perturbative treatment of the rotational effect is inadequate (see the last paragraph of this section). We shall adopt the so-called “traditional approximation” (Chapman & Lindzen 1970; Unno et al. 1989; Bildsten et al. 1996), where the Coriolis forces associated with the horizontal component of the spin angular velocity are neglected. This allows the mode eigenfunction to be separated into angular and radial components. Note that the traditional approximation is strictly valid only for high-order g-modes for which $\omega_r \ll N$ and $\xi_h \gg \xi_r$ (where $\xi_h \sim \xi_\theta \sim \xi_\phi$ is the horizontal Lagrangian displacement and ξ_r is the radial displacement), and for $\Omega_s \ll N$ (so that the radial component of the Coriolis force can be neglected compared to the buoyancy force). While these conditions are not valid everywhere inside the star (e.g., near the center ξ_r is of the same order as ξ_h, and $\omega_r, \Omega_s > N$; also, near the surface, $\omega_r, \Omega_s > N$), in the region $(0.6 \lesssim r/R \lesssim 0.9)$ where the mode propagates, they are reasonably satisfied (see below).

Neglecting the perturbation in gravitational potential (Cowling approximation), the radial Lagrangian displacement and Eulerian pressure perturbation can be written as

$$\xi_r(\mathbf{r}) = \xi_r(r) H_{jm}(\theta) e^{im\phi}, \quad \delta P(\mathbf{r}) = \delta P(r) H_{jm}(\theta) e^{im\phi}, \tag{4}$$

where $H_{jm}(\theta)$ is the Hough function (jm are the indices analogous to lm in the spherical harmonic Y_{lm}), satisfying the Laplace tidal equation (Chapman & Lindzen 1970; Bildsten et al. 1996):

$$\frac{\partial}{\partial\mu}\left(\frac{1-\mu^2}{1-q^2\mu^2}\frac{\partial H_{jm}}{\partial\mu}\right) - \frac{m^2 H_{jm}}{(1-\mu^2)(1-q^2\mu^2)} - \frac{qm(1+q^2\mu^2)H_{jm}}{(1-q^2\mu^2)^2} = -\lambda H_{jm}, \tag{5}$$

with $\mu = \cos\theta$ and $q = 2\Omega_s/\omega_r$. The eigenvalue, λ, depends on m and q. For $q \to 0$ (the nonrotating case), the function $H_{jm}(\theta)e^{im\phi}$ becomes $Y_{lm}(\theta,\phi)$ while λ degenerates into $l(l+1)$. The other eigenfunctions of the mode are related to δP and ξ_r via

$$\begin{aligned}
\delta\rho(\mathbf{r}) &= \delta\rho(r) H_{jm} e^{im\phi} = \left(\frac{\delta P}{c_s^2} + \frac{\rho N^2}{g}\xi_r\right) H_{jm} e^{im\phi}, && (6)\\
\xi_\theta(\mathbf{r}) &= \frac{\xi_\perp(r)}{1-q^2\mu^2}\left(\frac{\partial H_{jm}}{\partial\theta} + \frac{mq\mu}{\sin\theta}H_{jm}\right) e^{im\phi}, && (7)\\
\xi_\phi(\mathbf{r}) &= \frac{i\xi_\perp(r)}{1-q^2\mu^2}\left(q\mu\frac{\partial H_{jm}}{\partial\theta} + \frac{m}{\sin\theta}H_{jm}\right) e^{im\phi}, && (8)
\end{aligned}$$

where c_s is the sound speed and $\xi_\perp(r) = \delta P(r)/(\rho r \omega_r^2)$. Separating out the angular dependence, the fluid continuity equation and Euler equation (in the rotating frame) reduce to a set of coupled radial equations:

$$\begin{aligned}
\frac{d}{dr}\delta P &= -\frac{g}{c_s^2}\delta P + \rho(\omega_r^2 - N^2)\xi_r, && (9)\\
\frac{d}{dr}(r^2\xi_r) &= \frac{g}{c_s^2}(r^2\xi_r) - \frac{r^2}{c_s^2\rho}\delta P + \frac{\lambda}{\omega_r^2\rho}\delta P. && (10)
\end{aligned}$$

The properties of the Hough function have been extensively studied (Longuet-Higgins 1967). We have adopted the numerical approach of Bildsten et al. (1996) in our calculation. For given q and m, the eigenvalue λ is obtained by solving equation (5). To obtain the eigen-frequency ω_r, we solve the radial equations (9)-(10) subjected to the following boundary conditions: (i) At the surface $r = R$, we have the standard requirement $\Delta P = \delta P - \rho g \xi_r = 0$; (ii) At a small r close the the center, regularity requires $\delta P/\rho = Y_0 r^\beta$, $\xi_r = (\beta/\omega_r^2) Y_0 r^{\beta-1}$ (Y_0 is a constant), where β is determined from $\beta(1+\beta) = \lambda$, or $\beta = \left(\sqrt{1+4\lambda} - 1\right)/2$. Once ω_r is found, the actual rotation rate Ω_s is recovered from $\Omega_s = q\omega_r/2$.

Figure 2 shows the frequencies (in the inertial frame) of the $j = m = 2$ g_1 modes for model AU and UU as functions of the spin frequencies (These modes have $l = 2$ in the $\Omega_s = 0$ limit). Note that as Ω_s increases, ω_r increases while ω_i decreases.

An approximate solution the the radial equation can be obtained by the WKB analysis. Substituting $r^2\xi_r \propto e^{ikr}$ and $\delta P/\rho \propto e^{ikr}$ into the radial equations, we obtain the dispersion relation for g-modes:

$$k^2 \simeq \frac{\lambda}{\omega_r^2 r^2}(N^2 - \omega_r^2) \tag{11}$$



Figure 2. The frequencies of the $j = m = 2$ g$_1$ modes in the inertial frame for model AU and UU as functions of the spin frequency. The onset of instability occurs at $\nu_{i,\rm mode} = 0$. The solid lines are numerical results obtained by solving the radial equations; The dashed lines correspond to the WKB limits.



Figure 3. Density perturbation $\delta\rho$ (in arbitrary unit) as a function of radius for the $j = m = 2$ g$_1$ mode of model UU. The solid line is for $\nu_s = 0$, and the dashed line is for $\nu_s \simeq 140$ Hz (corresponding to $q = 1.3$).

This indicates that $\omega_r \propto \sqrt{\lambda}$. In fact, with the condition $\int_{r_1}^{r_2} k\,dr = (n + C)\pi$ (n is the number of radial node, C is a constant of order unity, r_1 and r_2 are the inner and outer turning points where $\omega_r = N$), we can write

$$\nu_r \simeq \frac{234\,\mathrm{Hz}}{n+C}\left(\frac{\lambda}{6}\right)^{1/2}\left(\frac{\int \nu_N\, d\ln r}{300\,\mathrm{Hz}}\right). \tag{12}$$

Thus we have, for the $j = 2$ modes, $\omega_r/\omega^{(0)} = \sqrt{\lambda/6}$ and $\Omega_s/\omega^{(0)} = q\sqrt{\lambda/24}$, where $\omega^{(0)}$ is the corresponding mode frequency at zero rotation. The WKB results are also plotted in Fig. 2, showing remarkable agreement with the actual numerical result. Clearly, $\omega_i = 0$ occurs at $q = 1$, which corresponds to $\lambda = 11.13$ from the solution of the $m = 2$ Laplace equation (eq. [5]). Thus, the onset of instability occurs when the ratio of spin frequency Ω_s and the nonrotating mode frequency $\omega^{(0)}$ satisfies

$$\frac{\Omega_s}{\omega^{(0)}} = \left(\frac{11.13}{24}\right)^{1/2} = 0.68 \tag{13}$$

The actual numerical calculations give slightly smaller critical spin frequency (see Fig. 2). For $\bar{\Omega}_s \equiv \Omega_s/\omega^{(0)} < 1$, in the WKB limit, the $j = m = 2$ mode frequency can be fitted to the following analytic expression to within 1%: [‡]

$$\frac{\omega_i}{\omega^{(0)}} = 1 - \frac{5}{3}\bar{\Omega}_s + \frac{13}{42}\bar{\Omega}_s^2 - 0.064\,\bar{\Omega}_s^{4.6}. \tag{14}$$

Figure 3 gives two examples of the radial eigenfunctions. We see that the mode is concentrated in the outer region of the stellar core ($0.6 \lesssim r/R \lesssim 0.9$), and rotation tends to "squeeze" the mode into that region. This is because rotation increases ω_r and narrows the region in which the wave can propagate ($\omega_r < N$). The dispacements ξ_r and $\xi_\perp$ can be quite significant near the surface, but the associated energy density (proportional to $\rho|\vec{\xi}|^2$) is significant only in the propagation zone.

We can similarly calculate the frequencies of higher-order g-modes. The results are even closer to the WKB limit (as expected). These modes tend to be unstable at lower spin frequencies. Similarly, the higher-m modes are also destablized by gravitational waves. We shall not be concerned with these higher-order modes since they have smaller growth rates due to gravitational radiation and higher damping rates due to viscosities as compared to the $m = 2$, g$_1$ mode (see Section 4).

Finally, we discuss the validity of the two approximations adopted in our calculations. As far as the mode frequency is concerned, the Cowling approximation is an excellent approximation for g-modes: For nonrotating stars, our mode frequencies agree with those of Lai (1994) (where the full equations are solved) to within 2%. The Cowling approximation does introduce an error (up to a factor of 80%) to the quadrupole moment of the mode (see §4). However, the uncertainty in the nuclear equation of state (particularly the nuclear symmetry energy which determines the g-mode properties) is presumably larger. As noted before, the conditions $\omega_r \ll N$ and $\Omega_s \ll N$, needed for the validity of the traditional approximation, are satisfied in the propagation zone of the mode, although they are clearly violated near the stellar center and surface. Our numerical results indicate that for the g$_1$ mode, the inequality $|\xi_\perp| > |\xi_r|$ is achieved only for $r/R \gtrsim 0.6 - 0.7$ (depending on the rotation rate), and $|\xi_\perp| \gg |\xi_r|$ is achieved only near the stellar surface [indeed $\xi_\perp/\xi_r = GM/(R^3\omega_r^2) \gg 1$ at $r = R$]. Thus the traditional approximation is not rigorously justified for the g$_1$ mode, although it becomes increasingly valid for higher-order modes. Clearly, without an exact, full calculation, which is currently beyond reach, it is difficult to know definitively how good the traditional approximation is when applying to the low-order modes. One way to assess the validity of our results is to consider the $\omega^{(0)} \gg \Omega_s$ limit using the linear theory, which gives, for the $l = m = 2$ mode:

[‡] A series solution (in q) of eq. (5) yields $\lambda = 6+2q+10q^2/7+\cdots$ for $j = m = 2$ (Bildsten & Ushomirsky 1996, private communication). In eq. (14), the linear and quadratic terms are derived from this analytic expansion of λ for small q and the last term is based on our numerical fitting.

$$\omega_i = \omega^{(0)} - 2(1 - C_{nl})\Omega_s, \tag{15}$$

where

$$C_{n2} = \frac{\int_0^R \rho\, r^2\, (2\xi_r\xi_\perp + \xi_\perp^2)\, dr}{\int_0^R \rho\, r^2\, [\xi_r^2 + 6\xi_\perp^2]\, dr} \tag{16}$$

(e.g., Unno et al. 1989). If we assume $|\xi_\perp| \gg |\xi_r|$, we find $C_{nl} = 1/6$ and eq. (15) reduces to the first two terms of eq. (14). For model UU, our numerical calculation yields $C_{nl} = 0.104,\ 0.132,\ 0.148,\ 0.155$ for g_1,g_2,g_3,g_4 respectively, corresponding to $2(1 - C_{nl}) = 1.79,\ 1.74,\ 1.70,\ 1.69$. These values are close to the corresponding value $(5/3 = 1.67)$ obtained from the traditional approximation. Since our calculation indicates that the instability occurs at $\Omega_s < \omega^{(0)}$ (see eq. [13]), we are confident that our qualitative conclusion regarding the secular instability of g-modes is firm; the critical ratio $\Omega_s/\omega^{(0)}$ for instability obtained in this paper (eq. [13]) is likely to be accurate to within $\sim 20\%$.

4 DRIVING AND DAMPING OF THE MODES

The condition that ω_i changes sign is only a necessary condition for the CFS instability. For the instability to be of interest, it must grow sufficiently fast and that the growth rate must overcome the possible damping rates due to viscosities. In this section we determine the mode growth rate and damping rate.

The timescale τ associated with any dissipative process is given by

$$\frac{1}{\tau} = -\frac{1}{2E}\frac{dE}{dt}, \tag{17}$$

where E is the canonical mode energy in the rotating frame (Friedman & Schutz 1978a):

$$\begin{aligned} E \;&=\; \frac{1}{2}\int d^3x \bigg[\rho\,\omega_r^2\,\vec{\xi}^* \cdot \vec{\xi} + \left(\frac{\delta P}{\rho} - \delta\Phi\right)\delta\rho^* \\ &\qquad + (\nabla\cdot\vec{\xi})\,\vec{\xi}^* \cdot (\nabla p - c_s^2\nabla\rho)\bigg]. \end{aligned} \tag{18}$$

We normalize the mode amplitude according to

$$\int d^3x\,\rho\,\vec{\xi}^* \cdot \vec{\xi} = \int_0^R dr\, r^2 \rho\left(\xi_r^2 + \Lambda\xi_\perp^2\right) = 1, \tag{19}$$

where Λ is evaluated using

$$\begin{aligned} \Lambda \;&=\; 2\pi\int_{-1}^{1}\frac{d\mu}{(1-q^2\mu^2)^2}\bigg[(1+q^2\mu^2)(1-\mu^2)\left(\frac{\partial H_{jm}}{\partial\mu}\right)^2 \\ &\qquad + \frac{m^2(1+q^2\mu^2)}{1-\mu^2}H_{jm}^2 - 4mq\mu H_{jm}\left(\frac{\partial H_{jm}}{\partial\mu}\right)\bigg], \end{aligned} \tag{20}$$

and the Hough function is normlized via $2\pi\int d\mu\, H_{jm}^2 = 1$. The g-mode energy is dominated by the first term in eq. (18), which gives

$$E \simeq \frac{1}{2}\omega_r^2(1 + \mathcal{E}), \tag{21}$$

with

$$\mathcal{E} = \frac{1}{\omega_r^2}\int dr\, r^2\frac{\delta P(r)}{\rho}\delta\rho(r). \tag{22}$$

The dissipation due to gravitational waves is calculated from the multipole radiation formula derived by Lindblom et al. (1998) using the formalism of Thorne (1980):

$$\left(\frac{dE}{dt}\right)_{\mathrm{gr}} = -\omega_r\sum_{l\geq 2} N_l\omega_i^{2l+1}(|\delta D_{lm}|^2 + |\delta J_{lm}|^2). \tag{23}$$

where N_l is the coupling constant:

$$N_l = \frac{4\pi G}{c^{2l+1}}\frac{(l+1)(l+2)}{l(l-1)[(2l+1)!!]^2}, \tag{24}$$

and the mass and current multipole moments of the perturbed fluid are given by:

$$\delta D_{lm} = \int d^3x\,\delta\rho\, r^l Y_{lm}^*, \tag{25}$$

$$\delta J_{lm} = \frac{2}{c}\sqrt{\frac{l}{l+1}}\int d^3x\, r^l(\rho i\omega_r\vec{\xi} + \delta\rho\mathbf{v})\cdot\vec{Y}_{lm}^{B*}, \tag{26}$$

and $\vec{Y}_{lm}^B$ is the magentic type vector spherical harmonic defined by Thorne (1980):

$$\vec{Y}_{lm}^B = \frac{1}{\sqrt{l(l+1)}}r\nabla\times(r\nabla Y_{lm}). \tag{27}$$

For the $j = m = 2$ mode, the dominant gravitational radiation comes from the mass quadrupole δD_{22}. We find

$$\frac{1}{\tau_{\mathrm{gr}}} \simeq \frac{25}{1+\mathcal{E}}\hat{\omega}_i^5\hat{\omega}_r^{-1}\left(\frac{\delta D_{22}}{10^{-4}}\right)^2 M_{1.4}^3 R_1^{-4}\ \mathrm{yr}^{-1}, \tag{28}$$

where $\hat{\omega} = \omega/(GM/R^3)^{1/2}$, δD_{22} is dimensionless (in units where $G = M = R = 1$), $M_{1.4} = M/(1.4\,M_\odot)$ and $R_1 = R/(10\,\mathrm{km})$. Clearly, τ_{gr} changes from positive (damping) for $\omega_i > 0$ to negative (driving) for $\omega_i < 0$ (where the mode is unstable for inviscid stars).

Figure 4 shows the damping/growth rate τ_{gr}^{-1} as a function of the stellar spin frequency ν_s for the g_1 mode of the two neutron star models. The change of τ_{gr}^{-1} with varying ν_s mainly comes from the changes in ω_i and ω_r, with the change in δD_{22} relatively minor. We see that when ν_s exceeds the critical spin rate for instability, the mode growth rate can be significantly larger than the damping rate for the nonrotating models (This is particularly evident for model AU).

The higher order mass multipole radiations (from δD_{l2}, with $l > 2$) are less important. One can show that the radiation from δJ_{22} identically vanishes. The leading order current multipole radiation comes from δJ_{32}, but it is much smaller than the mass quadrupole radiation.

The growth/damping timescale of g-modes due to gravitational radiation is typically much longer than that of f-modes (for which τ_{gr} is of order seconds). This comes about for two reasons: (i) G-modes have lower frequencies; (ii) The multipole moment δD_{lm} for g-modes is much smaller than that for the f-modes (for which δD_{lm} is of order unity). [§]

[§] The small δD_{lm} for g-modes can be understood from the sum rule (Reisenegger 1994) that must be satisfied for the multipole moments of all modes.



Figure 4. The gravitational radiation damping rate $\tau_{\rm gr}^{-1}$ as a function of spin frequency for the two g-modes depicted in Fig. 2. Positive $\tau_{\rm gr}^{-1}$ (for small ν_s) implies damping, while negative $\tau_{\rm gr}^{-1}$ implies driving of the mode by gravitational radiation. At $\nu_s = 0$, we have $\tau_{\rm gr}^{-1} \simeq 0.04$ yr^{-1} for model UU and $\tau_{\rm gr}^{-1} \simeq 6\times 10^{-5}$ yr^{-1} for model AU.

The dissipation of mode energy due to viscosities can be calculated from (Ipser & Lindblom 1991)

$$\left(\frac{dE}{dt}\right)_{\rm visc} = -\int d^3x \,\left(2\eta\,\delta\sigma^{ab}\delta\sigma^*_{ab} + \zeta|\delta\sigma|^2\right), \tag{29}$$

where $\delta\sigma_{ab}$ and $\delta\sigma$ are the shear and expansion of the perturbation, respectively:

$$\begin{aligned}\delta\sigma_{ab} &= \frac{i\omega_r}{2}\left(\nabla_a\xi_b + \nabla_b\xi_a - \frac{2}{3}\delta_{ab}\nabla_c\xi^c\right), && (30)\\ \delta\sigma &= i\omega_r\nabla_c\xi^c. && (31)\end{aligned}$$

The shear viscosity resulting from neutron-neutron scattering in the normal liquid core is given by (Flowers & Itoh 1979; Cutler & Lindblom 1987):

$$\eta = 1.1\times 10^{17} T_9^{-2}\left(\frac{\rho}{\rho_{\rm nuc}}\right)^{9/4}\ \ {\rm g/(cm\ s)}, \tag{32}$$

where T_9 is the temperature in units of 10^9 K, and $\rho_{\rm nuc} = 2.8\times 10^{14}$ g cm^{-3}. The bulk viscosity results from the tendency to achieve chemical equilibrium via the modified URCA process during the perturbation (Sawyer 1989):

$$\zeta = 4.8\times 10^{24} T_9^6 \left(\frac{\rho}{\rho_{\rm nuc}}\right)^2 \left(\frac{\omega_r}{1\ {\rm s}^{-1}}\right)^{-2}\ \ {\rm g/(cm\ s)}. \tag{33}$$

We have not been able to derive an explicit expression for $(dE/dt)_{\rm visc}$ from the g-mode wavefunctions to facilitate convenient numerical computation of the viscous damping rate. We shall therefore be contented with order of magnitude estimates. The mode damping rate due shear viscosity is approximately given by the rate at which momentum diffuses across a mode wavelength:

$$\frac{1}{\tau_{\rm shear}} \sim \frac{\eta}{\rho L^2} \simeq 10^{-2} L_1^{-2} T_9^{-2}\left(\frac{\rho}{\rho_{\rm nuc}}\right)^{5/4}\ \ {\rm yr}^{-1}, \tag{34}$$

where $L = 10 L_1$ km is the characteristic wavelength of the g-mode. Similarly, the mode damping rate due to bulk viscosity is estimated as

$$\frac{1}{\tau_{\rm bulk}} \sim 0.6\, L_1^{-2} T_9^6 \left(\frac{\rho}{\rho_{\rm nuc}}\right)\left(\frac{\nu_r}{150\ {\rm Hz}}\right)^{-2}\ \ {\rm yr}^{-1}, \tag{35}$$

(where ν_r is the mode frequency in the rotating frame). The length scale of the mode is somewhat less than 10 km, as comparison with the numerical results for nonrotating g-modes (Lai 1994) indicates that eq. (34) (with $L_1 = 1$) underestimates $1/\tau_{\rm shear}$ by a factor of ten, while eq. (35) underestimates $1/\tau_{\rm bulk}$ by a factor of three. When the star rotates faster than the mode frequency at zero rotation, the transverse wavelength of the mode decreases as $\sim 2R/q$ (for $q \gg 1$), and the radial wavelength also tends to be smaller (see Fig. 3). For the rotation rate of interest in this paper (Ω_s is at most a few times $\omega^{(0)}$), the mode wave function is only modestly changed by rotation, and we expect that rotation modifies the viscous damping rates by at most a factor of a few. Also note that the mode frequency (in the rotating frame) ν_r increases with increasing spin frequency (e.g., for model UU, we have $\nu_r = 296$ Hz when $\nu_s = 300$ Hz), and this tends to reduce our estimate for $\tau_{\rm bulk}^{-1}$. In any case, the uncertainty in our estimate of viscous damping rate is probably comparable to the uncertainly in our understanding of the microscopic viscosity of neutron star matter.

The total damping rate of the g-mode is given by

$$\frac{1}{\tau} = \frac{1}{\tau_{\rm gr}} + \frac{1}{\tau_{\rm shear}} + \frac{1}{\tau_{\rm bulk}}. \tag{36}$$

The mode is unstable when $\tau^{-1} < 0$. Despite the uncertainty in our estimates for $\tau_{\rm shear}$ and $\tau_{\rm bulk}$, by comparing Fig. 4 with eqs. (34) and (35) we can readily conclude that for $T \gtrsim 10^9$ K the g-mode instability is suppressed by the bulk viscosity, while for $T \lesssim 10^9$ K it is likely to be suppressed by the shear viscosity. For a narrow range of neutron star temperatures, around a few times 10^8 K to 10^9 K (depending on the detail of viscous dissipation), the g_1 mode may be unstable due to gravitational radiation. To achieve this instability, the rotation rate of the star must be at least a factor of two to three greater than the mode frequency at zero rotation (see Fig. 4).

So far we have only focused on the $j = m = 2$ g_1 mode. As the order of the g-modes increases, both the mode frequency and δD_{lm} decreases, which tends to reduce $|\tau_{\rm gr}^{-1}|$. Also, the viscous mode damping rate increases with increasing mode order. Therefore we expect that the CFS instabilities of higher-order g-modes are completely suppressed by viscous dissipations.

5 VISCOSITY DRIVEN INSTABILITY?

For an incompressible rotating Maclaurin spheroid, the $l = -m = 2$ f-mode becomes unstable due to viscosity at the same point as the CFS instability of the $l = m = 2$ mode (Chandrasekhar 1987). The $l = -m > 2$ viscous instability occurs at higher rotation rates. Recent studies indicate the instability occurs only for neutron stars with very stiff equation of state (Bonazzola et al. 1997).

The viscous instability sets in when the mode frequency ω_r in the rotating frame goes through zero. Figure 5 shows



Figure 5. The frequencies (in the rotating frame) of the $j = -m = 2$ g_1 modes for the two neutron star models (UU and AU) as functions of the spin frequency. For comparison, the dashed lines show the results for the $j = -m = 3$ g_1 modes.

the frequencies of the $j = -m = 2$ modes for the two neutron star models (UU and AU). For comparison, the frequencies of the $j = -m = 3$ modes are also shown. All these modes follow the approximate WKB scaling relation $\omega_r \propto \sqrt{\lambda}$ (see Sect. 3). We see that while ω_r is reduced by rotation, it never goes to zero, up to the maximum possible rotation rate.[¶] We therefore conclude that viscosity-driven instability is absent for g-modes.

6 DISCUSSION

We have shown in this paper that g-modes in an inviscid neutron star are susceptible to the CFS instability when the rotation rate is comparable to or greater than the mode frequencies. However, viscous dissipation tends to suppress the instability except possibly in a narrow range of temperatures around 10^9 K.

Comparing to f-modes, the g-mode instability has the advantage that it sets in at a lower rotation rate. Comparing to r-modes, the g-mode instability is much weaker. Calculations (Lindblom et al. 1998; Andersson et al. 1998a) indicate that at spin frequency of 300 Hz, the growth rate of r-mode is approximately 10^3 yr^{-1}, more than three orders of magnitude larger than $1/\tau_{\rm gr}$ for the g-modes studied in this paper. Thus for isolated neutron stars, the r-mode instability plays an much more important role than the g-mode in determining the spin evolution of the stars (Owen et al. 1998; Andersson et al. 1998b).

However, one should not dismiss possible importance of neutron star g-modes and their stabilities in other situations. In a merging neutron star binary, g-modes can be resonantly excited by the companion star, with the consequence of changing the orbital decay rate (Reisenegger & Goldreich 1994; Lai 1994). Rotation may significantly change the strength of the resonance (Lai 1997), giving rise to measurable effect in the gravitational wave form (Ho & Lai 1999). In another situation, when the neutron star is surrounded by a disk, tidal coupling between the stellar g-modes and disk density waves can drive the the unstable modes, which may affect the stellar rotation. By contrast, r-modes possess no mass quadrupole moment, and the coupling with disk is mediated only through higher order multipole moments and thus is expected to be weak. We wish to study some of these issues in a future paper.

ACKNOWLEDGMENTS

Part of this paper was completed while the author was visiting the Aspen Centre for Physics during May-June of 1998. He thanks the referee for constructive comments which improved this paper, and acknowledges support from a NASA ATP grant, and a research fellowship from the Alfred P. Sloan Foundation.

REFERENCES

Andersson, N., 1998, ApJ, 502, 708
Andersson, N., Kokkotas, K.D., Schutz, B.F., 1998a, ApJ, 510, 846
Andersson, N., Kokkotas, K.D., Stergioulas, N., 1998b, preprint astro-ph/9806089.
Baym, G., Bethe, H.A, & Pethick, C.J. 1971, Nucl. Phys. A175, 225
Bildsten, L. 1998, ApJ, 501, L89
Bildsten, L., & Cumming, A. 1998, ApJ, 506, 842
Bildsten, L., Cutler, C., 1995, ApJ, 449, 800
Bildsten, L., Ushomirsky, G., Cutler, C., 1996, ApJ, 460, 827
Bonazzola, S., Frieben, J., Gourgoulhon, E., 1998, A&A, 331, 280
Chandrasekhar, S., 1970, Phys. Rev. Lett., 24, 611
Chandrasekhar, S., 1987, Ellipsoidal Figures of Equilibrium (Dover Pub.: New York)
Chapman, S., Lindzen, R.S., 1970, Atmospheric Tides (Gordon and Breach: New York)
Cutler, C., Lindblom, L., 1987, ApJ, 314, 234
Cutler, C., Lindblom, L., 1992, ApJ, 385, 630
Finn, L.S., 1987, MNRAS, 227, 265
Flowers, E., Itoh, N., 1979, ApJ, 230, 847.
Friedman, J.L., Morsink, S.M., 1998, ApJ, 502, 714
Friedman, J.L., Schutz, B.F., 1978a, ApJ, 221, 937
Friedman, J.L., Schutz, B.F., 1978b, ApJ, 222, 281
Ho, W.C.G., Lai, D. 1999, MNRAS, in press (astro-ph/9812116)
Ipser, J.R., Lindblom, L., 1990, ApJ, 355, 226
Ipser, J.R., Lindblom, L., 1991, ApJ, 373, 213
Lai, D., 1994, MNRAS, 270, 611
Lai, D., 1997, ApJ, 490, 847
Levin, Y. 1998, astro-ph/9810471
Lindblom, L., 1995, ApJ, 438, 265
Lindblom, L., Owen, B.J., Morsink, S.M., 1998, Phys. Rev. Lett., 80, 4843
Longuet-Higgins, M.S., 1967, Phil. Trans. R. Soc. London A, 262, 511
McDermott, P.N., & Taam, R. E. 1987, ApJ, 318, 278
McDermott, P.N., Van Horn, H.M., Hansen, C.J., 1988, ApJ, 325, 725
Owen, B.J., Lindblom, L., Cutler, C., Schutz, B.F., Vecchio, A., Andersson, N., 1998, Phys. Rev. D 5808, 4020

[¶] Note that as the rotation rate approaches the maximum value, our calculation of the mode frequency breaks down since the rotational distortion becomes significant.

Papaloizou, J., Pringle, J.E. 1978, MNRAS, 182, 423

Provost, J., Berthomieu, G., Rocca, A., 1981, A&A, 94, 126

Reisenegger, A, 1994, ApJ, 432, 296

Reisenegger, A, Goldreich, P., 1992, ApJ, 395, 240

Reisenegger, A, Goldreich, P., 1994, ApJ, 426, 688

Saio, H., 1982, ApJ, 256, 717

Stergioulas, N., & Friedman, J.L., 1998, ApJ, 492, 301

Strohmayer, T. E., 1993, ApJ, 417, 273.

Strohmayer, T. E., & Lee, U. 1996, ApJ, 467, 773

Thorne, K.S., 1980, Rev. Mod. Phys., 52, 299

Unno, W. et al., 1989, Nonradial Oscillation of Stars (Univ. of Tokyo Press)

Wiringa, R.B., Fiks, V., Fabrocini, A., 1988, Phys. Rev. C, 38, 1010

Yoshida, S., Eriguchi, Y., 1995, ApJ, 438, 830